arXiv:math-ph/0702059v1 16 Feb 2007

# THE CONSERVATION OF MASS-MOMENT PARAMETERS

DAN COMĂNESCU

Abstract. In this paper we study a concept of mass-moment parameter which is the generalization of *the mass* and *the moments of inertia* of a continuous media. We shall present some interesting kinematical results in the hypothesis that a set of mass-moment parameters are conserved in a motion of a continuous media.

*Mathematics Subject Classification: 74A05; 70B10; 70S10*

*Keywords: continuum mechanics; kinematics; mass-moment parameter.*

## 1. Introduction

A mass-moment parameter of a continuous media is a kinematical parameter which is used to describe the distribution of the matter in the continuous media. The most common mass-moment parameters are *the mass* and *the moments of inertia*. The mass-moment parameters, with the exception of *the mass*, are used especially in the theory of rigid bodies. In a motion of a rigid body the moments of inertia are conserved with respect to a frame of reference rigidly connected to the body.

The objective of our study are:

- to present the concept of mass-moment parameter which is used in this paper;
- to introduce the conservation of a mass-moment parameter for a motion of a continuous media;
- to deduce the local laws in the cases in which one or two mass-moment parameters are conserved in a motion;
- to study the conservation of some moments of inertia.

## 2. The mass-moment parameters

In this paper we consider the mass-moment parameters of the form (with respect to a frame of reference $\mathcal{R}$):

$$P(\mathcal{P},t)=\int_{P_t}\rho(\vec{x},t)p(\vec{x},t)dv \tag{2.1}$$

where $t$ is a time-moment, $\mathcal{P}$ is a part of the continuous media, $P_t$ is the image of $\mathcal{P}$ (at time $t$), $\vec{x}$ is the position vector of a particle (at time $t$), with respect to frame of reference $\mathcal{R}$, $\rho$ is *the mass density* and $p$ is the *reduced density* of the mass moment parameter $P$.

The mass-moment parameter $P$ is defined by a Lebesgue integral with respect to the volume measure. The functions $\rho$ and $p$ are supposed to be continuous.

For the reduced density $p \equiv 1$ on obtain *the mass* of the continuous media:

$$M(\mathcal{P},t) = \int_{P_t} \rho(\vec{x},t)dv \tag{2.2}$$

The *moment of inertia with respect to the point* $Q$ is defined by the integral:

$$I_Q(\mathcal{P},t) = \int_{P_t} \rho(\vec{x},t)d^2(\vec{x},\vec{x}_Q)dv \tag{2.3}$$

where $d(\vec{x},\vec{x}_Q)$ is the distance between an arbitrary point of $P_t$ and $Q$.
The *moment of inertia with respect to the plane* $\Pi$ is defined by the integral:

$$I_\Pi(\mathcal{P},t) = \int_{P_t} \rho(\vec{x},t)d^2(\vec{x},\Pi)dv \tag{2.4}$$

where $d(\vec{x},\Pi)$ is the distance between an arbitrary point of $P_t$ and the plane $\Pi$.
The *moment of inertia with respect to the* $\Delta$*-axis* is defined by the integral:

$$I_\Delta(\mathcal{P},t) = \int_{P_t} \rho(\vec{x},t)d^2(\vec{x},\Delta)dv \tag{2.5}$$

where $d(\vec{x},\Delta)$ is the distance between an arbitrary point of $P_t$ and the line $\Delta$.

## 3. The conservation of a set of mass-moment parameters

In this section we present some results concerning the conservation of one or two mass-moment parameters in a motion of a continuous media.
First, we deduce the local laws for the conservation of a mass-moment parameter.
Second, two conservation type results are deduced.

**Definition 3.1.** *A mass-moment parameter $P$ is conserved in a motion of the continuous media if we have:*

$$P(\mathcal{P},t_1) = P(\mathcal{P},t_2) \tag{3.1}$$

*for all parts $\mathcal{P}$ of the continuous media and for all two time-moments $t_1$ and $t_2$.*

We denote by $\vec{X}$ and $\vec{x}$ the position occupied by a particle at the initial moment and in the configuration at $t$. The law (3.1) has the form:

$$\int_{P_t} \rho(\vec{x},t)p(\vec{x},t)dv = \int_{P_0} \rho(\vec{X},0)p(\vec{X},0)dV \tag{3.2}$$

**Remark 3.1.** The problem of conservation of a mass-moment parameter is connected with the mathematical theory of integral invariants.

**Theorem 3.1.** ***(local law in material variables)*** *Let the $C^1$-map $\vec{x} = \chi(\vec{X},t)$ of the motion and $P$ a mass-moment parameter with $p$ the reduced density. The proposition are equivalents:*
*(i) $P$ is conserved in the motion;*
*(ii) for all time-moments $t$ and $\vec{X} \in D_0$ we have:*

$$\rho(\vec{x},t)p(\vec{x},t)J(\vec{X},t) = \rho(\vec{X},0)p(\vec{X},0) \tag{3.3}$$

*where* $D_0$ *is the image of the continuous media at the initial moment and* $J$ *the determinant of the deformation gradient; i.e.:*

$$J(\vec{X},t) = det(\frac{\partial x_i}{\partial X_j}(\vec{X},t))_{i,j\in\{1,2,3\}}$$

*Proof.* We need the following result:

**Lemma 3.2.** *(see [8] pp. 47) If* $f : \mathcal{D} \subset \mathbb{R}^3 \to \mathbb{R}$ *is a continuous function on the domain* $\mathcal{D}$ *such that for all domains* $D \subset \mathcal{D}$ *is satisfied* $\int_D f dv = 0$ *then* $f \equiv 0$ *.*

Using the relation (3.2) and the transformation of the integrals in the material variables we deduce that for all domains $P_0 \subset D_0$ and for all times $t$:

$$\int_{P_0} \rho(\vec{x},t)p(\vec{x},t)J(\vec{X},t) - \rho(\vec{X},0)p(\vec{X},0)dV = 0. \tag{3.4}$$

The theorem follows from (3.4) and Lemma 3.2. □

**Theorem 3.3.** ***(local law in spatial variables)*** *Let the* $C^2$*-map* $\vec{x} = \chi(\vec{X},t)$ *of the motion and* $P$ *a mass-moment parameter with the reduced density* $p$ *a* $C^1$*-function. The proposition are equivalents:*
*(i)* $P$ *is conserved in the motion;*
*(ii) for all time-moments* $t$ *and* $\vec{x} \in D_t$ *we have:*

$$\frac{d(\rho p)}{dt}(\vec{x},t) + (\rho p)(\vec{x},t)div_x\vec{v}(\vec{x},t) = 0 \tag{3.5}$$

*where* $\vec{v}$ *is the velocity and* $div_x\vec{v}$ *is the divergence of* $\vec{v}$ *with respect to the spatial variables.*

*Proof.* We need the Euler's lemma (see [8],pp. 36).

**Lemma 3.4.** ***(Euler)*** *For a* $C^1$*-motion* $\vec{x} = \chi(\vec{X},t)$ *of a continuous media is satisfied the relation:*

$$\frac{dJ}{dt}(\vec{X},t) = J(\vec{X},t)div_x\vec{v}(\vec{x},t).\text{''}$$

We have the sequence of equivalences:

$$(i) \;\Leftrightarrow\; \frac{d}{dt}\int_{P_t} \rho p \, dv = 0 \;\; \forall \mathcal{P} \;\Leftrightarrow\; \int_{P_0} \frac{dJ}{dt}\rho p + \frac{d\,\rho p}{dt} J \, dV = 0 \;\; \forall \mathcal{P}$$

Using Euler's lemma and Lemma 3.2 we deduce:

$$(i) \;\Leftrightarrow\; \int_{P_t} \frac{d(\rho p)}{dt} + (\rho p)div_x(\vec{v},t)\, dv = 0 \;\; \forall \mathcal{P} \;\Leftrightarrow\; (ii)$$

□

We shall study now a case in which *two* mass-moment parameters are conserved.

**Theorem 3.5.** *Let a $C^1$-motion of a continuous media $\vec{x} = \chi(\vec{X}, t)$. We consider two mass-media parameters $P_1$ and $P_2$ with reduced densities $p_1$ and $p_2$ which are conserved in the motion. Then, for all particles and time-moments it is satisfied:*

$$p_1(\vec{X}, 0)p_2(\vec{x}, t) = p_1(\vec{x}, t)p_2(\vec{X}, 0) \tag{3.6}$$

*Proof.* Following Theorem 3.1 one obtains:

$$\rho(\vec{x}, t)p_1(\vec{x}, t)J(\vec{X}, t) = \rho(\vec{X}, 0)p_1(\vec{X}, 0)$$

$$\rho(\vec{x}, t)p_2(\vec{x}, t)J(\vec{X}, t) = \rho(\vec{X}, 0)p_2(\vec{X}, 0)$$

Using the properties $\rho > 0$ and $J \neq 0$ we deduce our result. □

**Theorem 3.6.** *If, moreover, $p_2 \neq 0$ we deduce:*

$$\frac{p_1}{p_2}(\vec{x}, t) = \frac{p_1}{p_2}(\vec{X}, 0) \tag{3.7}$$

**Theorem 3.7.** *Let a $C^1$-motion of a continuous media $\vec{x} = \chi(\vec{X}, t)$ and $P$ a mass-moment parameter with $p$ the reduced density.*
*If the mass $M$ (see (2.2)) and $P$ are conserved in the motion then for all particles and time-moments we have:*

$$p(\vec{x}, t) = p(\vec{X}, 0) \tag{3.8}$$

## 4. Applications

In this section are presented some applications of the results of section 3.

4.1. **Conservation of mass.** If the mass $M$ (see (2.2)) is conserved in a $C^1$-motion $\vec{x} = \chi(\vec{X}, t)$ of a continuous media then, using Theorem 3.1, is obtained *the material equation of continuity* (Euler 1762):

$$\rho(\vec{X}, 0) = \rho(\vec{x}, t)J(\vec{X}, t) \tag{4.1}$$

If the motion is a $C^2$-function and $\rho$ is a $C^1$-function a consequence of conservation of mass is (see Theorem 3.3) *the spatial equation of continuity* (Euler 1757):

$$\frac{d\rho}{dt}(\vec{x}, t) + \rho(\vec{x}, t)div_x\vec{v}(\vec{x}, t) = 0 \tag{4.2}$$

4.2. **Conservation of mass and of an other mass-moment parameter.** In this paragraph we suppose that a continuous media has a $C^1$-motion such that the mass $M$ (see (2.2)) is conserved.

Using Theorem 3.7 is easy to obtain the following results.

**Theorem 4.1.** *Let $O$ a fixed point. The following are equivalents:*
*(i) the moment of inertia with respect to the point $O$ is conserved in the motion;*
*(ii) the motion of an arbitrary particle is on a sphere with the center $O$.*

**Theorem 4.2.** *Let* $\Pi$ *a fixed plane. The following are equivalents:*
*(i) the moment of inertia with respect to the plane* $\Pi$ *is conserved in the motion;*
*(ii) the motion of an arbitrary particle is in a plane parallel with* $\Pi$.

**Theorem 4.3.** *Let* $\Delta$ *a fixed axis. The following are equivalents:*
*(i) the moment of inertia with respect to the* $\Delta$*-axis is conserved in the motion;*
*(ii) the motion of an arbitrary particle is on a circular cylinder with* $\Delta$ *as the axis of symmetry .*

4.3. **A condition for the equilibrium of a continuous media.** Let a $C^1$-motion of a continuous media with respect to a spatial frame of reference $Ox_1x_2x_3$. We suppose that the mass $M$ is conserved in the motion. We denote by $I_O$ the moment of inertia with respect to the point $O$, $I_{x_i}$ $(i \in \{1,2,3\})$ the moment of inertia with respect to the $Ox_i$-axis, $I_{Ox_ix_j}$ $(i,j \in \{1,2,3\})$ the moment of inertia with respect to the plane $Ox_ix_j$. We have the relations (see [14] pp. 612-613):

$$I_O = \frac{1}{2}(I_{x_1} + I_{x_2} + I_{x_3}) \tag{4.3}$$

$$I_O = I_{Ox_1x_2} + I_{Ox_2x_3} + I_{Ox_3x_1} \tag{4.4}$$

$$I_O = I_{x_i} + I_{Ox_jx_k}, \ \ \{i,j,k\} = \{1,2,3\} \tag{4.5}$$

$$I_{x_i} = I_{Ox_ix_j} + I_{Ox_ix_k}, \ \ \{i,j,k\} = \{1,2,3\} \tag{4.6}$$

$$I_{Ox_ix_j} = \frac{1}{2}(I_{x_i} + I_{x_j} - I_{x_k}), \ \ \{i,j,k\} = \{1,2,3\} \tag{4.7}$$

**Theorem 4.4.** *If three of seven mass-moment parameters from the set* $I_O, I_{x_1}, I_{x_2}, I_{x_3}, I_{Ox_1x_2}, I_{Ox_2x_3}, I_{Ox_3x_1}$ *are conserved in the motion then the continuous media is in an equilibrium state.*

*Proof.* Using the relations (4.3)-(4.7) we deduce that all the seven mass-moment parameters are conserved in the motion of the continuous media. Applying the Theorem 3.6 for the mass-moment parameters $I_{Ox_1x_2}, I_{Ox_2x_3}, I_{Ox_3x_1}$ and we obtain:

$$\begin{cases} x_1^2(\vec{X},t) = X_1^2 \\ x_2^2(\vec{X},t) = X_2^2 \\ x_3^2(\vec{X},t) = X_3^2 \end{cases}$$

The functions $t \to x_i(\vec{X},t)$ are continuous with the initial conditions $x_i(\vec{x},t) = X_i$. The result is straightforward. $\square$

## References

[1] ]cite.Arnold741V.I. Arnold, *Mathematical methods of classical mechanics*, Nauka, Moscow, 1974.

[2] ]cite.Balint982St. Balint, *Lecţii de mecanică teoretică. Mecanica solidului rigid*, Tip. Univ. de Vest, Timişoara, 1998.

[3] ]cite.Balint963______, *Lecţii de mecanică teoretică. Mecanica mediilor continue*, Tip. Univ. de Vest, Timişoara, 1996.

[4] ]cite.Camenschi004G. Camenschi, *Introducere în mecanica mediilor continue deformabile*, Ed. Univ. Bucureşti, Bucureşti, 2000.

[5] ]cite.Comanescu045D. Comănescu, *Modele şi metode în mecanica punctului material*, Mirton, Timişoara, 2004.

[6] ]cite.Dragos036L. Dragoş, *Mathematical methods in Aerodynamics*, Kluwer Academic Pub. and Ed. Academiei Române, Bucureşti, 2003.

[7] ]cite.Dragos997______, *Mecanica fluidelor, vol. I. Teoria generală. Fluidul ideal incompresibil*, Ed. Academiei Române, Bucureşti, 1999.

[8] ]cite.Dragos838______, *Principiile mecanicii mediilor continue*, Ed. Tehnică, Bucureşti, 1983.

[9] ]cite.Dragos769______, *Principiile mecanicii analitice*, Ed. Tehnică, Bucureşti, 1976.

[10] ]cite.Iacob8910C. Iacob, *Matematică aplicată şi mecanică*, Ed. Academiei R.S.R., Bucureşti, 1989.

[11] ]cite.Iacob8011______, *Mecanică teoretică*, Ed. Didactică şi Pedagogică, Bucureşti, 1980.

[12] ]cite.Soos8312E. Soós and C. Teodosiu, *Calcul tensorial cu aplicaţii în mecanica solidelor*, Ed. Ştiinţifică şi Enciclopedică, Bucureşti, 1983.

[13] ]cite.Truesdell7413C. Truesdell, *Introduction a la Mecanique des Milieux Continus*, Mason, Paris, 1974.

[14] ]cite.Valcovici6814V. Vâlcovici, St Bălan, and R. Voinea, *Mecanică teoretică, ed. a III-a*, Ed. Tehnică, Bucureşti, 1968.

ADDRESS OF AUTHOR:, DEPARTMENT OF MATHEMARTICS, WEST UNIVERSITY OF TIMIŞOARA, BD. PÂRVAN NR. 4, TIMIŞOARA, ROMANIA
*E-mail address*: `comanescu@math.uvt.ro`